




# Marbocen 100 mg/ml

**Injectable solution**  
**Marbofloxacin 100 mg/ml**

- *Respiratory complex*
- *MMA Syndrome*
- *Mastitis*



**Short withdrawal periods**



# Marbocen 100 mg/ml

## Injectable solution Marbofloxacin 100 mg/ml

### COMPOSITION PER ML:

#### Active substance:

Marbofloxacin ..... 100,0 mg

Other excipients, q.s.

### TARGET SPECIES INDICATIONS:

#### Cattle:

- Treatment of respiratory infections caused by of *Histophilus somni*, *Mannheimia haemolytica*, *Mycoplasma bovis* and *Pasteurella multocida* sensitive to marbofloxacin.
- Treatment of acute mastitis caused by strains of *E. coli* sensitive to marbofloxacin during the lactating period.

#### Pigs:

- Treatment of Postpartum Dysgalactia Syndrome - PPSD - (Metritis Mastitis Agalactia Syndrome), caused by bacterial strains susceptible to marbofloxacin.
- Treatment of respiratory infections in fattening pigs - The recommended dose is 2 mg marbofloxacin/ kg/day, (1 ml/50 kg BW) in a single daily intramuscular injection for 3 to 5 days.

### DOSAGE FOR EACH SPECIES, ROUTES AND METHOD OF ADMINISTRATION:

#### Cattle:

**Respiratory infections:** The recommended dosage is 8 mg marbofloxacin/kg body weight (2 ml product/25 kg BW) in a single injection by intramuscular route. If the volume to be injected is more than 20 ml, it should be divided between two or more injection sites.

In cases of respiratory infections caused by *Mycoplasma bovis*, the recommended dose is 2 mg marbofloxacin/kg body weight (1 ml product/50 kg BW), in a single daily injection for 3 to 5 consecutive days, by intramuscular or subcutaneous route. The first injection may be given by the intravenous route.

**Acute mastitis:** The recommended dosage is 2 mg marbofloxacin/kg body weight (1 ml product/50 kg BW) in a single daily injection for 3 consecutive days by intramuscular or subcutaneous route. The first injection may also be given by the intravenous route.

#### Porcine:

#### Treatment of Postpartum Dysgalactia Syndrome (PPDS) - Metritis Mastitis Agalactia Syndrome:

The recommended dose is 2 mg marbofloxacin/kg body weight (1 ml/50 kg bw) in a single daily injection by intramuscular via during 3 days.

#### Treatment of respiratory infections in fattening pigs:

The recommended dose is 2 mg marbofloxacin/ kg/day, (1 ml/50 kg BW) in a single daily intramuscular injection for 3 to 5 days.

### WITHDRAWAL PERIOD:

#### Bovine:

8 mg/kg in a single injection (IM): Meat: 3 days. Milk: 72 hours.

2 mg/kg during 3 or 5 days (IV/SC/IM): Meat: 6 days. Milk: 36 hours.

#### Porcine:

Meat: 4 days.

### PRESENTATIONS:

Vials of 100 ml and 250 ml.

### VETERINARY USE