


# Cendox<sup>500</sup>

**Doxycycline hyclate 500 mg/gr**



- **High solubility**
- **Prolonged stability**
- **Withdrawal time in pigs: 48 h**



# Cendox<sup>500</sup>

**Doxycycline hyclate 500 mg/gr**

**Powder for oral solution**

## COMPOSITION:

### Active substance:

Doxycycline (hyclate) ..... 500 mg

### Excipient:

Citric acid anhydrous q.s.....1 gr

## TARGET SPECIES AND INDICATIONS FOR USE:

**Pigs:** Treatment of infectious processes included in the Porcine Respiratory Disease Complex produced by microorganisms sensitive to doxycycline as are: *Pasteurella multocida* and *Mycoplasma hyopneumoniae*.

**Broilers:** treatment of colibacillosis and Chronic Respiratory Disease (CRD) caused by *Escherichia coli* and *Mycoplasma gallisepticum*.

**Pre-ruminant calves:** infections of the respiratory tract caused by microorganisms sensitive to doxycycline as are: *Pasteurella multocida* and *Mannheimia haemolytica*.

## AMOUNTS TO BE ADMINISTERED AND ADMINISTRATION ROUTE

**Poultry and pigs:** Oral via in the drinking water.

**Poultry:** 15 mg of doxycycline/kg b.w./day for 3-5 days (equivalent to 30 mg of product/kg b.w./ day).

**Pigs:** 10 mg of doxycycline/kg b.w./day for 5 days (equivalent to a 0,2 g of product/10 kg b.w./day).

**Pre-ruminant calves:** Oral via in milk at a temperature about 37°C and administered in the next 60 minutes. 10 mg of doxycycline / kg b.w./ day (equivalent to 0,2 g of product/10 kg b.w./ day) for 5 days.

Assess daily the necessary quantity of drug in accordance to the animal weight and add to the milk.

## WITHDRAWAL TIME:

**Meat:**

**Broilers:** 7 days

**Pigs:** 2 days

**Pre-ruminant calves:** 7 days

Not authorized for lactating animals whose milk is intended for human consumption, nor laying hens whose eggs are intended for human consumption.

## PACK SIZES

200 g and 1 kg