



corticen 2 mg/ml

**Injectable solution
Dexamethasone**

- *Maximum effectiveness*
- *Excellent syringability*
- *Wide spectrum of use*



- Inflammatory and allergic processes
- Induction of parturition in cattle
- Primary ketosis (acetonemia) in cattle
- Arthritis, bursitis or tenosynovitis in horses



corticen 2 mg/ml

Injectable solution Dexamethasone

COMPOSITION PER ML:

Active substance: Dexamethasone as Dexamethasone Sodium Phosphate.....2.0 mg

Excipients: Benzyl alcohol (E 1519).....15.6 mg

TARGET SPECIES:

Horses, cattle, pigs, dogs and cats.

INDICATIONS FOR USE:

Horses, cattle, pigs, dogs and cats: Treatment of inflammatory or allergic processes.

Cattle: Induction of labour and treatment of primary ketosis.

Horses: Treatment of arthritis, bursitis or synovitis.

DOSAGE FOR EACH SPECIES, ROUTES AND METHOD OF ADMINISTRATION:

Administration routes: Horses: IV, IM or intra-articular use. **Cattle, pigs, dogs and cats:** IM use.

Treatment of inflammatory or allergic processes:

Horses, cattle, pigs: 1.5 ml / 50 kg.

Dogs, cats: 0.5 ml / 10 kg.

For the treatment of primary ketosis in cattle (acetonemia): 5-10 ml / 500 kg b.w., depending on the duration of signs, is recommended. Care should be taken not to overdose in Jersey, Guernsey and/or related breeds. Higher doses will be required if signs have been present for a prolonged period of time or if treating animals that have relapsed. In most cases with recent symptoms, a cure will be obtained with a single dose.

For induction of labour: 10 ml / 500 kg bw as a single dose after day 260 of gestation. Calving will usually occur within 48 to 72 hours.

For the treatment of arthritis, bursitis or tenosynovitis, by intra-articular injection, in horses: Dose 1 ml - 5 ml. These quantities are not specific and are quoted for reference only. Injections into intra-articular spaces or bursae should be preceded by the removal of an equivalent volume of synovial fluid.

WITHDRAWAL PERIODS:

Cattle: Meat and offal: 8 days. Milk: 72 hours

Pigs: Meat and offal: 2 days.

Horses: Meat and offal: 8 days. Not authorised for use in horses producing milk for human consumption.

PRESENTATION:

Vials of 100 ml. Clinical packages containing 10 vials of 100 ml.

VETERINARY USE