




Cenflox 200 mg/ml

Solution for use in drinking water
Enrofloxacin 200 mg/ml



ENROFLOXACIN



High concentration

Multispecies

High profitability



CenfloX 200 mg/ml

Solution for use in drinking water

Enrofloxacin 200 mg/ml



COMPOSITION PER ML:

Enrofloxacin.....200 mg
Excipients, q.s.

TARGET SPECIES:

Chickens, turkeys and rabbits.

INDICATIONS:

Chickens: Treatment of infections caused by the following bacteria susceptible to enrofloxacin: *Mycoplasma gallisepticum*, *Mycoplasma synoviae*, *Avibacterium paragallinarum*, *Pasteurella multocida*.

Turkeys: Treatment of infections caused by the following bacteria susceptible to enrofloxacin: *Mycoplasma gallisepticum*, *Mycoplasma synoviae*, *Pasteurella multocida*.

Rabbits: For the treatment infectious diseases due to *Pasteurella multocida* and bacterial enteritis due to infection with *E.coli*.

DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION:

Administer in the drinking water.

The dilution should be prepared daily before its administration.

Chickens and turkeys: 10 mg enrofloxacin/kg bodyweight, equivalent to 0.05 ml veterinary medicinal product/kg bodyweight, per day for 3-5 consecutive days.

Rabbits: 10 mg of enrofloxacin/kg b.w., equivalent to 0.05 ml veterinary medicinal product/kg b.w., per day for 5 consecutive days.

WITHDRAWAL PERIOD:

Chickens: Meat and offal: 7 days

Turkeys: Meat and offal: 13 days

Rabbits: Meat and offal: 3 days

Not authorised for use in birds producing eggs for human consumption.

Do not administer to layer replacement birds within 14 days of coming into lay

PRESENTATION:

1L and 5L

VETERINARY USE