


Colicen 4.000.000 U.I./ml

Colistin (sulfate) 4.000.000 U.I./ml



**Treatment of gastrointestinal
problems produced by E. Coli and
Salmonella**



WITHDRAWAL PERIOD

- **Calves, lambs and pigs: Meat and offal: 1 day**
- **Chickens and turkeys: 1 day**
- **Egg: 0 days**



Colicen

4.000.000 U.I./ml



Colistin (sulfate) 4.000.000 U.I./ml

COMPOSITION PER ML:

Colistin (as sulfate) 4000000 IU

Excipients, q.s.

TARGET SPECIES:

Cattle (calves), sheep (lambs), pigs, chickens and turkeys.

POSOLOGY AND ADMINISTRATION VIA:

In drinking water/milk use:

Calves, lambs, pigs: 100 000 IU of colistin per kg body weight daily for 3-5 consecutive days in drinking water or milk (replacer) in calves, equivalent to 0.25 ml of the concentrate solution per 10 kg body weight per day for 3-5 days.

Chickens and turkeys: 75 000 IU of colistin per kg body weight daily for 3-5 consecutive days in drinking water, equivalent to 18.75 ml of the concentrate solution per Ton of body weight per day for 3-5 days.

Duration of treatment should be limited to the minimum time necessary for the treatment of the disease.

Any medicated water which is not consumed within 24 hours should be discarded.

Any medicated milk which is not consumed within 3 hours should be discarded.

Direct oral administration to individual animals: The recommended daily dose should be divided into two if the product is to be administered directly into the mouth of the animal. Prior to direct oral administration, the product should be diluted with a volume of drinking water equivalent to 2 x the volume of product concentrate to be administered.

WITHDRAWAL PERIOD:

Calves, lambs and pigs: Meat and offal: 1 day.

Chickens and turkeys: Meat and offal: 1 day. Egg: 0 days.

PRESENTATION:

1 L and 5 L

VETERINARY USE