



daxton


Injectable solution

Tulathromycin 100 mg/ml

**New generation
antibiotic**

**Low dose and
cost of treatment**

One Shot



- **Respiratory disease in bovine and swine (BRD/SRD)**
- **Sheep foot rot**



daxton

Injectable solution

Tulathromycin 100 mg/ml

COMPOSITION PER ML:

Tulathromycin100 mg
Excipients, q.s.

TARGET SPECIES:

Cattle, pigs and sheep.

INDICATIONS FOR USE:

Cattle: Treatment and metaphylaxis of bovine respiratory disease (BRD) associated with *Mannheimia haemolytica*, *Pasteurella multocida*, *Histophilus somni* and *Mycoplasma bovis* susceptible to tulathromycin. Treatment of infectious bovine keratoconjunctivitis (IBK) associated with *Moraxella bovis* susceptible to tulathromycin.

Pigs: Treatment and metaphylaxis of swine respiratory disease (SRD) associated with *Actinobacillus pleuropneumoniae*, *Pasteurella multocida*, *Mycoplasma hyopneumoniae*, *Haemophilus parasuis* and *Bordetella bronchiseptica* susceptible to tulathromycin.

Sheep: Treatment of the early stages of infectious pododermatitis (foot rot) associated with virulent *Dichelobacter nodosus* requiring systemic treatment.

DOSAGE AND ADMINISTRATION VIA:

Cattle: Subcutaneous use. A single subcutaneous injection of 1 ml/40 kg b.w.

For treatment of cattle over 300 kg b.w., divide the dose so that no more than 7.5 ml are injected at one site.

Pigs: Intramuscular use. A single intramuscular injection of 1 ml/40 kg b.w.

For treatment of pigs over 80 kg b.w., divide the dose so that no more than 2 ml are injected at one site.

Sheep: Intramuscular use. A single intramuscular injection 1 ml/40 kg b.w.

WITHDRAWAL PERIODS:

Meat and offal

Cattle: 22 days.

Pigs: 13 days.

Sheep: 16 days.

Milk: Not authorised for use in animals producing milk for human consumption.

Do not use in pregnant animals, which are intended to produce milk for human consumption, within 2 months of expected parturition.

SHELF LIFE:

Shelf life of the veterinary medicinal product as packaged for sale: 3 years.

Shelf life after first opening the immediate packaging: 28 days.

PRESENTATIONS:

Vials of 50 ml, 100 ml and 250 ml.

VETERINARY USE