

Florfenicen 300 mg/ml

Florfenicol

Injectable solution for bovine and pigs



**Straight
to the infection**



Florfenicen 300 mg/ml

Injectable solution for bovine and pigs



COMPOSITION:

Each ml contains:

Florfenicol..... 300 mg/ml

Excipients, n.q.f.

PROPERTIES:

Florfenicol acts by inhibiting protein synthesis of bacteria to be bacteriostatic ribosomal level. Laboratory tests have shown that the florfenicol is active against pathogens commonly isolated in bovine respiratory disease, including those caused by *Mannheimia haemolytica*, *Pasteurella multocida* and *H. somni*, and including porcine respiratory disease caused by *Pasteurella multocida* and *Actinobacillus pleuropneumoniae*.

Florfenicol is considered bacteriostatic, but in vitro studies demonstrate bactericidal activity against *Mannheimia haemolytica*, *Pasteurella multocida*, *Actinobacillus pleuropneumoniae* and *H. somni*.

TARGET SPECIES:

Bovine and pigs.

INDICATIONS:

Bovine: treatment of respiratory tract infections caused by strains of *Mannheimia haemolytica*, *Pasteurella multocida* and *H. somni* sensitive to florfenicol.

Pigs: Treatment of acute outbreaks of respiratory disease caused by strains of *Actinobacillus pleuropneumoniae* and *Pasteurella multocida* susceptible to florfenicol.

DOSAGE AND ADMINISTRATION:

Route of administration: intramuscular or subcutaneous injection in bovine, intramuscular in pigs.

Bovine: IM via: 20 mg / kg body weight (1 ml/15 kg) twice with an interval of 48 hours using a 16-gauge needle.

SC via: 40 mg / kg body weight (2 ml/15 kg) single dose using a 16-gauge needle. The administered by injection volume should not exceed 10 ml in both routes of administration (intramuscular and subcutaneous). The injection should be given in the neck.

Pigs: 15 mg / kg body weight (1 ml / 20 kg) intramuscularly in the neck muscle twice with an interval of 48 hours using a 16-gauge needle. The administered by injection volume should not exceed 3 ml.

WITHDRAWAL TIME:

Bovine: **Meat:** Intramuscular (20 mg / kg body weight, twice): 30 days. Subcutaneously (40 mg / kg body weight, single dose): 44 days.

Milk: Its use is not authorized in lactating animals producing milk for human consumption.

Pigs: **Meat and offal:** 18 days.

PRESENTATION:

Vials of 100 ml and 250 ml.

VETERINARY USE



C. Dels Boters, s/n. • Pol. Ind. Mas Ferrer • 43205 REUS (SPAIN) • Tel. 34 977 757 273 • info.export@cenavisa.com

cenavisa.com