Florfenicen 300 mg/ml

Injectable solution
Florfenicol 300 mg/ml

Straight to the infection

cenavisa
ANIMAL HEALTH

Florfenicen 300 mg/ml

Injectable solution
Florfenicol 300 mg/ml

Straight to the infection

COMPOSITION PER ML:
Florfenicol . . . . . 300 mg
Excipients, n.q.f.

TARGET SPECIES:
Cattle, sheep and pigs.

INDICATIONS:
Cattle: Diseases caused by florfenicol susceptible bacteria: therapeutic treatment of respiratory tract infections in cattle due to *Mannheimia haemolytica, Pasteurella multocida* and *Hisophilus somni.*
Sheep: Diseases caused by florfenicol susceptible bacteria: therapeutic treatment of respiratory tract infections in sheep due to *Mannheimia haemolytica* and *Pasteurella multocida.*
Pigs: Treatment of acute outbreaks of respiratory disease caused by strains of *Actinobacillus pleuropneumoniae* and *Pasteurella multocida susceptible to florfenicol.*

DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION:
Route of administration: intramuscular and subcutaneous for cattle; intramuscular for sheep and pigs.
Cattle: IM route: 20 mg/kg bodyweight (1ml/15kg) to be administered twice 48 hours apart using a 16 gauge needle. **SC route:** 40 mg/kg bodyweight (2ml/15kg) to be administered once only using a 16 gauge needle. The dose volume given at any one injection site should not exceed 10 ml for both routes of administration (intramuscular and subcutaneous) The injection should only be given in the neck.
Ovine: 20 mg/kg bodyweight (1 ml per 15 kg) per day by intramuscular injection during three consecutive days. The dose volume given at any one injection site should not exceed 4 ml.
Pigs: 15 mg/kg bodyweight (1 ml per 20 kg) by intramuscular injection into the neck muscle twice at 48 hour intervals using a 16-gauge needle. The volume administered per injection site should not exceed 3 ml.

WITHDRAWAL PERIOD:
Cattle: Meat: by IM (at 20 mg/kg bodyweight, twice): 30 days, by SC (at 40 mg/kg bodyweight, once): 44 days.
Milk: Not authorized for use in lactating animals producing milk for human consumption including pregnant animals intended to produce milk for human consumption.
Sheep: Meat: 39 days.
Milk: Not authorized for use in lactating animals producing milk for human consumption including pregnant animals intended to produce milk for human consumption.
Pigs: Meat: 18 days.

PRESENTATIONS:
Vials of 100 and 250 ml

VETERINARY USE

cenavisa ANIMAL HEALTH
C. dels Boters, 4 • Pol. Ind. Mas Ferrer • 43205 REUS (SPAIN) • Tel. +34 977 757 273
info.export@cenavisa.com