

Ketoprocen

Ketoprocen 100 mg/ml

Injectable solution for horses, cattle and pigs



- ***Pain***
- ***Fever***
- ***Inflammation***



WITHDRAWAL PERIODS

Horses: Meat: 4 days. Milk: zero hours.

Cattle: Meat: 4 days. Milk: zero hours.

Pigs: Meat: 4 days.



Ketoprofen

Injectable solution

COMPOSITION:

Active substance: Ketoprofen.....100 mg/ml

Excipient: Benzyl alcohol..... 10 mg/ml



- **Pain**
- **Fever**
- **Inflammation**

TARGET SPECIES:

Equine, cattle and swine.

INDICATIONS:

Horse:

Diseases affecting the osteoarticular and muscular-skeletal system associated with acute pain and inflammation: Lameness of traumatic origin Arthritis, Osteitis, spavin, Tendinitis, bursitis, Naviculitis, Laminitis, Myositis, Ketoprofen is also indicated for post-surgical inflammation, symptomatic therapy of colic and fever.

Cattle:

Diseases associated with inflammation, pain or fever: Respiratory diseases, Mastitis, Osteoarticular and muscular-skeletal disorders such as lameness, arthritis and to ease uprise post parturition, Injuries.

Swine:

Diseases associated with inflammation, pain or fever: Treatment associated with the Mastitis Metritis Agalactia (MMA) Syndrome, Respiratory tract infections, Symptomatic treatment of fever,

For the short-term relief of post operative pain associated with minor soft tissue surgery such as castration in piglets.

- Where necessary ketoprofen should be combined with appropriate antimicrobial therapy.

POSOLGY:

Horse: 2.2 mg ketoprofen/kg body weight/day intravenously once daily for up to 3 to 5 consecutive days, i.e. 1 ml per 45 kg body weight.

In order to treat colic one injection is normally sufficient. A second administration of ketoprofen requires a reassessment of the patient's clinical status.

Cattle: 3 mg ketoprofen/kg body weight/day intravenously or deep intramuscularly once daily for up to 3 consecutive days, i.e. 3 ml per 100 kg body weight.

Swine: 3 mg ketoprofen/kg body weight as a single deep intramuscular injection, i.e. 3 ml per 100 kg body weight (= 0.03 ml/kg).

For reduction of post-operative pain the product should be injected 10 - 30 minutes before surgical intervention. Particular care should be taken with regard to the accuracy of dosing including the use of an appropriate dosing device (i.e.: low dose syringe) and proper determination of body weight.

PACK SIZES:

Vials of 100 ml and 250 ml.

VETERINARY USE



C. Dels Boters, s/n. • Pol. Ind. Mas Ferrer • 43205 REUS (SPAIN) • Tel. 34 977 757 273 • info.export@cenavisa.com

cenavisa.com