



ketoprofen 100 mg/ml

Injectable solution

Ketoprofen 100 mg/ml



- ***Pain***
- ***Fever***
- ***Inflammation***



Withdrawal periods

Horses: Meat: 4 days. Milk: 0 hours.

Cattle: Meat: 4 days. Milk: 0 hours.

Pigs: Meat: 4 days.



ketoprofen 100 mg/ml

Injectable solution Ketoprofen 100 mg/ml

COMPOSITION PER ML:

Ketoprofen100 mg/ml
Benzyl alcohol10 mg/ml
Excipients, q.s

TARGET SPECIES:

Horses, cattle and pigs.

INDICATIONS:

Horses:

- Alleviation of inflammation and pain associated with musculoskeletal disorders.
- Symptomatic treatment of visceral pain associated with colic. Postoperative pain and inflammation.

Cattle:

- Anti-inflammatory and analgesic treatment of diseases of musculoskeletal system.
Reducing the pyrexia and distress associated with bacterial respiratory disease when used in connection with anti-microbial therapy as appropriate.
- Reducing mammary oedema.
- Anti-inflammatory, analgesic and anti-pyretic treatment in acute clinical mastitis signs of mastitis in conjunction with antimicrobial therapy.

Pigs:

- Reducing the pyrexia in respiratory diseases.
- Treatment of Postpartum Dysgalactia Syndrome, Mastitis Metritis Agalactia Syndrome.

DOSAGE FOR EACH SPECIES, ROUTE(S) AND METHOD OF ADMINISTRATION:

Intramuscular or intravenous use.

Horses: Intravenous injection.

- For use in musculoskeletal conditions, the dosage is 2.2 mg of ketoprofen per kg b.w., equivalent to 1 ml of product for each 45 kg of bodyweight, once daily for up to 3 to 5 days.
- For use in colic, the dosage is 2.2 mg of ketoprofen per kg, i.e. 1 ml of product for each 45 kg of bodyweight, in a single injection. Generally a single injection is sufficient: any additional injection should be preceded by a clinical re-assessment of the animal.

Cattle: Intramuscular or intravenous injection. Recommended dose: 3 mg ketoprofen/kg b.w., equivalent to 1 ml of product for each 33 kg of bodyweight, once daily for up to 1 to 3 days.

Pigs: Intramuscular injection. Recommended dose: 3 mg/kg b.w., equivalent to 1 ml of product for each 33 kg of bodyweight, in a single dose.

WITHDRAWAL PERIODS:

Horses: Meat: 4 days. Milk: 0 hours.

Cattle: Meat: 4 days. Milk: 0 hours.

Pig: Meat: 4 days.

PRESENTATION:

Vials of 100 ml and 250 ml.

Clinical packages containing 10 vials of 100 ml and 10 vials of 250 ml.

VETERINARY USE



- **Pain**
- **Fever**
- **Inflammation**

MORE INFORMATION

